

# MMP14/MT1-MMP Rabbit mAb

Catalog No.: A0067 **Recombinant** **3 Publications**

## Basic Information

**Observed MW**

52kDa/60kDa

**Calculated MW**

66kDa

**Category**

Primary antibody

**Applications**

WB, IHC-P, FC, ELISA

**Cross-Reactivity**

Human, Mouse, Rat

**CloneNo number**

ARC0211

## Background

Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. However, the protein encoded by this gene is a member of the membrane-type MMP (MT-MMP) subfamily; each member of this subfamily contains a potential transmembrane domain suggesting that these proteins are expressed at the cell surface rather than secreted. This protein activates MMP2 protein, and this activity may be involved in tumor invasion.

## Recommended Dilutions

**WB** 1:500 - 1:2000**IHC-P** 1:50 - 1:200**FC** 1:100 - 1:500

**ELISA** Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## Immunogen Information

**Gene ID**

4323

**Swiss Prot**

P50281

**Immunogen**

Synthetic peptide. This information is considered to be commercially sensitive.

**Synonyms**

MMP-14; MMP-X1; MT-MMP; MT1MMP; MTMMP1; WNCHRS; MT1-MMP; MT-MMP 1; MMP14/MT1-MMP

## Contact

 | 400-999-6126 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn) | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

**Source**

Rabbit

**Isotype**

IgG

**Purification**


Affinity purification

**Storage**


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.


Validation Data




Western blot analysis of various lysates using MMP14/MMP14/MT1-MMP Rabbit mAb (A0067) at 1:1000 dilution.  
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 90s.




Western blot analysis of various lysates using MMP14/MMP14/MT1-MMP Rabbit mAb (A0067) at 1:1000 dilution.  
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 10s.



Flow cytometry: 1X10<sup>6</sup> MCF7 cells (negative control,left) and HT-1080 (right) cells were intracellularly-stained with MMP14/MT1-MMP Rabbit mAb (A0067,2 µg/mL,orange line) or ABflo® 488 Rabbit IgG isotype control (AC042,2 µg/mL,blue line), followed by Alexa Fluor® 488 conjugated goat anti-rabbit pAb staining. Non-fluorescently stained cells were used as blank control (red line).




Flow cytometry: 1X10<sup>6</sup> MCF7 cells (negative control,left) and HT-1080 (right) cells were intracellularly-stained with MMP14/MT1-MMP Rabbit mAb (A0067,2 µg/mL,orange line) or ABflo® 488 Rabbit IgG isotype control (AC042,2 µg/mL,blue line), followed by Alexa Fluor® 488 conjugated goat anti-rabbit pAb staining. Non-fluorescently stained cells were used as blank control (red line).




Flow cytometry: 1X10<sup>6</sup> MCF7 cells (negative control,left) and HT-1080 (right) cells were intracellularly-stained with MMP14/MT1-MMP Rabbit mAb (A0067,2 µg/mL,orange line) or ABflo® 647 Rabbit IgG isotype control (AC042,2 µg/mL,blue line), followed by Alexa Fluor® 647 conjugated goat anti-rabbit pAb staining. Non-fluorescently stained cells were used as blank control (red line).


## Validation Data




Flow cytometry:  $1 \times 10^6$  MCF7 cells (negative control, left) and HT-1080 (right) cells were intracellularly-stained with MMP14/MT1-MMP Rabbit mAb (A0067, 2  $\mu\text{g/mL}$ , orange line) or ABflo® 647 Rabbit IgG isotype control (AC042, 2  $\mu\text{g/mL}$ , blue line), followed by Alexa Fluor® 647 conjugated goat anti-rabbit pAb staining. Non-fluorescently stained cells were used as blank control (red line).



Immunohistochemistry analysis of paraffin-embedded Human liver tissue using MMP14/MT1-MMP Rabbit mAb (A0067) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human spleen tissue using MMP14/MT1-MMP Rabbit mAb (A0067) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse kidney tissue using MMP14/MT1-MMP Rabbit mAb (A0067) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.