

# ALDH1A1 Rabbit mAb

Catalog No.: A0157

Recombinant

4 Publications

## Basic Information

**Observed MW**

55kDa

**Calculated MW**

55kDa

**Category**

Primary antibody

**Applications**

WB,IF/ICC,IP,ELISA

**Cross-Reactivity**

Human, Mouse, Rat

**CloneNo number**

ARC52440

## Background

The protein encoded by this gene belongs to the aldehyde dehydrogenase family. Aldehyde dehydrogenase is the next enzyme after alcohol dehydrogenase in the major pathway of alcohol metabolism. There are two major aldehyde dehydrogenase isozymes in the liver, cytosolic and mitochondrial, which are encoded by distinct genes, and can be distinguished by their electrophoretic mobility, kinetic properties, and subcellular localization. This gene encodes the cytosolic isozyme. Studies in mice show that through its role in retinol metabolism, this gene may also be involved in the regulation of the metabolic responses to high-fat diet.

## Recommended Dilutions

**WB** 1:10000 - 1:60000**IF/ICC** 1:8000 - 1:32000**IP** 0.5µg-4µg antibody for  
200µg-400µg extracts of  
whole cells**ELISA** Recommended starting  
concentration is 1 µg/mL.  
Please optimize the  
concentration based on  
your specific assay  
requirements.

## Contact

 | 400-999-6126 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn) | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Immunogen Information

**Gene ID**

216

**Swiss Prot**

P00352

**Immunogen**

A synthetic peptide corresponding to a sequence within amino acids 250-350 of human ALDH1A1 (NP\_000680.2).

**Synonyms**

ALDC; ALDH1; HEL-9; HEL12; PUMB1; ALDH11; RALDH1; ALDH-E1; HEL-S-53e; ALDH1A1

## Product Information

**Source**

Rabbit

**Isotype**

IgG

**Purification**


Affinity purification

**Storage**


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.09% Sodium azide, 0.05% BSA, 50% glycerol, pH7.3.


Validation Data



Immunoprecipitation analysis of 300 µg extracts of A-549 cells using 3 µg ALDH1A1 Rabbit mAb antibody (A0157). Western blot was performed from the immunoprecipitate using ALDH1A1 Rabbit mAb (A0157) at a dilution of 1:20000.




Western blot analysis of various lysates, using ALDH1A1 Rabbit mAb (A0157) at 1:20000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Negative control (NC): THP-1. Exposure time: 10s.




Western blot analysis of various lysates, using ALDH1A1 Rabbit mAb (A0157) at 1:20000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 90s.

## Validation Data



Confocal imaging of Hep G2 cells using ALDH1A1 Rabbit mAb (A0157, dilution 1:8000) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with  $\alpha$ -Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.



Confocal imaging of A549 cells using ALDH1A1 Rabbit mAb (A0157, dilution 1:8000) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with  $\alpha$ -Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.