

CDKN1A/p21CIP1 Rabbit pAb

Catalog No.: A11454

5 Publications

Basic Information

Observed MW

Refer to figures

Calculated MW

18kDa

Category

Primary antibody

Applications

IHC-P, ELISA

Cross-Reactivity

Mouse, Rat

Background

This gene encodes a potent cyclin-dependent kinase inhibitor. The encoded protein binds to and inhibits the activity of cyclin-cyclin-dependent kinase2 or -cyclin-dependent kinase4 complexes, and thus functions as a regulator of cell cycle progression at G1. The expression of this gene is tightly controlled by the tumor suppressor protein p53, through which this protein mediates the p53-dependent cell cycle G1 phase arrest in response to a variety of stress stimuli. This protein can interact with proliferating cell nuclear antigen, a DNA polymerase accessory factor, and plays a regulatory role in S phase DNA replication and DNA damage repair. This protein was reported to be specifically cleaved by CASP3-like caspases, which thus leads to a dramatic activation of cyclin-dependent kinase2, and may be instrumental in the execution of apoptosis following caspase activation. Mice that lack this gene have the ability to regenerate damaged or missing tissue. Multiple alternatively spliced variants have been found for this gene.

Recommended Dilutions

IHC-P 1:50 - 1:200**ELISA** Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

1026

Swiss Prot

P38936

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

P21; CIP1; SDI1; WAF1; CAP20; CDKN1; MDA-6; p21CIP1; CDKN1A/p21CIP1

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Validation Data



Immunohistochemistry analysis of paraffin-embedded Mouse liver tissue using CDKN1A/p21CIP1 Rabbit pAb (A11454) at a dilution of 1:100 (40x lens). Microwave antigen retrieval was performed with 0.01 M PBS buffer (pH 7.2±0.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse brain tissue using CDKN1A/p21CIP1 Rabbit pAb (A11454) at a dilution of 1:100 (40x lens). Microwave antigen retrieval was performed with 0.01 M PBS buffer (pH 7.2±0.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat brain tissue using CDKN1A/p21CIP1 Rabbit pAb (A11454) at a dilution of 1:100 (40x lens). Microwave antigen retrieval was performed with 0.01 M PBS buffer (pH 7.2±0.2) prior to IHC staining.