# **TBCB Rabbit pAb**

Catalog No.: A13248 5 Publications



### **Basic Information**

#### **Observed MW**

37kDa

### **Calculated MW**

27kDa

### Category

Primary antibody

### **Applications**

WB,IHC-P,ELISA

#### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

Predicted to be involved in cell differentiation and nervous system development. Located in cytosol.

# **Recommended Dilutions**

**WB** 1:1000 - 1:2000

**IHC-P** 1:50 - 1:200

Recommended starting concentration is 1 µg/mL.

Please optimize the concentration based on your specific assay requirements.

# **Immunogen Information**

**Gene ID Swiss Prot** 1155 Q99426

#### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

### **Synonyms**

CG22; CKAP1; CKAPI; TBCB

## **Contact**

<u>a</u>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
$\overline{a}$	ı	www.ahclonal.com.cn


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### **Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.




Western blot analysis of various lysates using TBCB Rabbit pAb (A13248) at 1:3000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit lgG (H+L) (AS014) at 1:10000 dilution.


Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020).


Exposure time: 30s.



Immunohistochemistry analysis of paraffinembedded Rat brain using TBCB Rabbit pAb (A13248) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffinembedded Human placenta using TBCB Rabbit pAb (A13248) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffinembedded Mouse kidney using TBCB Rabbit pAb (A13248) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.