# FOXA1 Rabbit pAb

Catalog No.: A15278 9 Publications



# **Basic Information**

### **Observed MW**

50kDa

### **Calculated MW**

49kDa

### Category

Primary antibody

### **Applications**

WB,IHC-P,ELISA

#### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

This gene encodes a member of the forkhead class of DNA-binding proteins. These hepatocyte nuclear factors are transcriptional activators for liver-specific transcripts such as albumin and transthyretin, and they also interact with chromatin. Similar family members in mice have roles in the regulation of metabolism and in the differentiation of the pancreas and liver.

# **Recommended Dilutions**

WB 1:500 - 1:1000

**IHC-P** 1:50 - 1:200

**ELISA** Recommended starting concentration is 1  $\mu$ g/mL.

Please optimize the concentration based on your specific assay requirements.

# **Immunogen Information**

**Gene ID**3169

Swiss Prot
P55317

#### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

## **Synonyms**

HNF3A; TCF3A; FOXA1

# **Contact**

<b>a</b>	400-999-6126
×	cn.market@abclonal.com.cn
$\overline{a}$	www.abclonal.com.cn


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.




Western blot analysis of lysates from Rat testis, using FOXA1 Rabbit pAb (A15278) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit  $lgG\ (H+L)\ (AS014)$  at 1:10000 dilution.


Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020).


Exposure time: 30s.




Immunohistochemistry analysis of paraffinembedded Rat lung using FOXA1 Rabbit pAb (A15278) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffinembedded Mouse stomach using FOXA1 Rabbit pAb (A15278) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffinembedded Human colon carcinoma using FOXA1 Rabbit pAb (A15278) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffinembedded Human breast cancer using FOXA1 Rabbit pAb (A15278) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.