

[KO Validated] Ki67 Rabbit mAb

Catalog No.: A20018 **KO Validated** **Recombinant** **85 Publications**

Basic Information

Observed MW

359 kDa and below

Calculated MW

359 kDa

Category

Primary antibody

Applications

WB,IF/ICC,IF-F,IHC-P,mIHC,ELISA

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC5050-01

Background

Enables protein C-terminus binding activity. Involved in regulation of chromosome segregation and regulation of mitotic nuclear division. Located in chromosome; nuclear body; and nucleolus. Colocalizes with condensed chromosome. Implicated in Crohn's disease; breast cancer; human immunodeficiency virus infectious disease; and pancreatic cancer. Biomarker of several diseases, including Barrett's esophagus; autoimmune disease of musculoskeletal system (multiple); endocrine gland cancer (multiple); gastrointestinal system cancer (multiple); and interstitial cystitis.

Recommended Dilutions

WB	1:2000 - 1:10000
IF/ICC	1:50 - 1:200
IF-F	1:200 - 1:800
IHC-P	1:1000 - 1:4000
mIHC	1:500 - 1:4000
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Immunogen Information

Gene ID

4288

Swiss Prot

P46013

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

KIA; MIB-; MIB-1; PPP1R105; Ki67

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Validation Data




Western blot analysis of lysates from wild type (WT) and Ki67 knockout (KO) HeLa cells using [KO Validated] Ki67 Rabbit mAb (A20018) at 1:5000 dilution incubated overnight at 4°C. HeLa cells were treated with Nocadazole (1 µg/mL) at 37°C for 16 hours
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
 Lysates/proteins: 25 µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Basic Kit (RM00020).
 Exposure time: 45 s.




Confocal imaging of HeLa cells using [KO Validated] Ki67 Rabbit mAb (A20018, dilution 1:100)(Red) followed by a further incubation with Cy3-conjugated Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α -Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.




Immunohistochemistry analysis of paraffin-embedded Human breast cancer tissue using [KO Validated] Ki67 Rabbit mAb (A20018) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human colon carcinoma tissue using [KO Validated] Ki67 Rabbit mAb (A20018) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human tonsil tissue using [KO Validated] Ki67 Rabbit mAb (A20018) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



The multiplex IHC analysis on paraffin-embedded Human breast cancer tissue using the following specific primary antibodies and tyramide signal amplification (TSA) reagents (RK05903) : Ki67 Rabbit mAb (A20018, 1:500) with TSA-TYR-520 (Green), HER2/ErbB2 Rabbit mAb (A21248, 1:200) with TSA-TYR-570 (Red), and Androgen Receptor Rabbit mAb (A19611, 1:400) with



Confocal imaging of frozen sections of Rat liver tissue using [KO Validated] Ki67 Rabbit mAb (A20018, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Microwave antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.

Validation Data

TSA-TYR-690 (Magenta), DAPI (Blue) was used for nuclear staining. Prior to multiplex IHC staining, high-pressure antigen retrieval was performed using 0.01M citrate buffer at pH 6.0. The analysis was completed using a 20x objective lens.