

TriMethyl-Histone H3-K27 Rabbit mAb

Catalog No.: A22006 **Recombinant** **6 Publications**

Basic Information

Observed MW

17 kDa

Calculated MW

15 kDa

Category

Primary antibody

Applications

WB,IP,IF/ICC,IHC-P,DB,ChIP,CUT&Tag,ELISA

Cross-Reactivity

Human, Mouse, Rat, Other (Wide Range Predicted)

CloneNo number

ARC54169

Recommended Dilutions

WB	1:10000 - 1:40000
DB	1:2000 - 1:20000
IHC-P	1:500 - 1:3000
IF/ICC	1:200 - 1:1000
IP	0.5µg-4µg antibody for 200µg-600µg extracts of whole cells
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
ChIP	5µg antibody for 5µg-10µg of Chromatin
CUT&Tag	10 ⁵ cells /1 µg

Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.

Immunogen Information

Gene ID

8290/8350

Swiss Prot

Q16695/P68431

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

H3t; H3.4; H3/g; H3FT; H3C16; HIST3H3; TriMethyl-Histone H3-K27

Product Information

Source

Rabbit

Isotype

IgG

Purification

Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Contact

 | 400-999-6126


 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn


Validation Data




Immunoprecipitation of TriMethyl-Histone H3-K27 from 600 µg extracts of 293F cells was performed using 5 µg of TriMethyl-Histone H3-K27 Rabbit mAb (A22006). Rabbit IgG isotype control (AC005) was used to precipitate the Control IgG sample. IP samples were eluted with 1X reducing Laemmli Buffer. The Input lane represents 10% of the total input. Western blot analysis of immunoprecipitates was conducted using TriMethyl-Histone H3-K27 Rabbit mAb (A22006) at a dilution of 1:20000.




Western blot analysis of various lysates using TriMethyl-Histone H3-K27 Rabbit mAb (A22006) at 1:11000 dilution incubated overnight at 4°C. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Negative control (NC): H3 protein. Exposure time: 30s.




Chromatin immunoprecipitation analysis of extracts of HeLa cells, using TriMethyl-Histone H3-K27 antibody (A22006) and rabbit IgG. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.



Confocal imaging of HeLa cells using TriMethyl-Histone H3-K27 Rabbit mAb (A22006, dilution 1:200) followed by a




Confocal imaging of NIH/3T3 cells using TriMethyl-Histone H3-K27 Rabbit mAb (A22006, dilution 1:200) followed by a




Confocal imaging of C6 cells using TriMethyl-Histone H3-K27 Rabbit mAb (A22006, dilution 1:200) followed by a further

Validation Data


further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α -Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.




further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α -Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.




incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α -Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.




Immunohistochemistry analysis of paraffin-embedded Human liver tissue using TriMethyl-Histone H3-K27 Rabbit mAb (A22006) at a dilution of 1:700 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse spleen tissue using TriMethyl-Histone H3-K27 Rabbit mAb (A22006) at a dilution of 1:700 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat lung tissue using TriMethyl-Histone H3-K27 Rabbit mAb (A22006) at a dilution of 1:700 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



CUT&Tag was performed using the CUT&Tag Assay Kit(pAG-Tn5) for Illumina (RK20265) from 10^5 HeLa cells with 1 μ g Tri-Methyl-Histone H3-K27 Rabbit mAb(A22006), along with a Goat Anti-Rabbit IgG(H+L). The CUT&Tag results indicate the enrichment pattern of H3K27me3 in representative gene loci(MYT1).

CUT&Tag was performed using the CUT&Tag Assay Kit(pAG-Tn5) for Illumina (RK20265) from 10^5 HeLa cells with 1 μ g Tri-Methyl-Histone H3-K27 Rabbit mAb(A22006), along with a Goat Anti-Rabbit IgG(H+L). The CUT&Tag results indicate the enrichment pattern of H3K27me3 in representative gene loci(MYT1).

Dot-blot analysis of all sorts of peptides using TriMethyl-Histone H3-K27 Rabbit mAb (A22006) at 1:11000 dilution.