

APC Rabbit anti-Human ICAM-1/CD54 mAb

Catalog No.: A26917

Basic Information

Observed MW**Calculated MW**

58kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Human

CloneNo number

ARC54998

Conjugate

APC. Ex:650nm. Em:660nm.

Background

Intercellular cell adhesion molecule-1 (CD54 or ICAM-1) is a cell surface glycoprotein that belongs to the immunoglobulin superfamily (IgSF) of adhesion molecules. CD54 is expressed at low levels in diverse cell types, and is induced by cytokines (TNF- α , interleukin-1) and bacterial lipopolysaccharide. Apical localization of CD54 on endothelial cells (or basolateral localization on epithelial cells) is a prerequisite for leukocyte trafficking through the endothelial (or epithelial) barrier. Apical expression of CD54 on epithelial cells mediates pathogen invasion as well as host defense, a pattern also observed in tumors. CD54 also functions as a co-stimulator on antigen presenting cells, binding to its receptor LFA-1 (leukocyte function-associated antigen-1) on the surface of T cells during antigen presentation. Cross-linking of CD54 or binding to its ligand triggers activation of Src family kinases and the Rho/ROCK pathway. Phosphorylation on Tyr485 of CD54 is required for its association with SHP-2. SHP-2 seems essential for CD54-induced Src activation.

Recommended Dilutions

FC 5 μ l per 10^6 cells in
100 μ l volume

Immunogen Information

Gene ID

3383

Swiss Prot

P05362

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

BB2; CD54; P3.58

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

Validation Data



Flow cytometry: 1×10^6 Human PBMC were surface-stained with APC Rabbit anti-Human ICAM-1/CD54 mAb (A26917, 5 μ l/Test, orange line) or APC Rabbit IgG isotype control (A24173, 5 μ l/Test, blue line). Non-fluorescently stained cells were used as blank control (red line).



Flow cytometry: 1×10^6 Human PBMC were surface-stained with APC Rabbit IgG isotype control (A24173, 5 μ l/Test, left) or APC Rabbit anti-Human ICAM-1/CD54 mAb (A26917, 5 μ l/Test, right)