

Biotin Rabbit anti-Mouse/Rat CD31/PECAM1 mAb

Catalog No.: A27363

Basic Information

Observed MW

Calculated MW

69kDa/77kDa/80kDa/81kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Mouse, Rat

CloneNo number

ARC61018

Conjugate

Biotin

Background

Predicted to enable protein homodimerization activity; protein phosphatase binding activity; and transmembrane signaling receptor activity. Acts upstream of or within several processes, including Rho protein signal transduction; endothelial cell morphogenesis; and positive regulation of tyrosine phosphorylation of STAT protein. Located in several cellular components, including cell-cell contact zone; external side of plasma membrane; and smooth muscle contractile fiber. Is expressed in several structures, including alimentary system; cardiovascular system; central nervous system; extraembryonic component; and genitourinary system. Human ortholog(s) of this gene implicated in coronary artery disease (multiple); lung non-small cell carcinoma; neuroblastoma; and psoriatic arthritis. Orthologous to human PECAM1 (platelet and endothelial cell adhesion molecule 1).

Recommended Dilutions

FC ≤0.25 µg per million cells
in 100 µl volume

Immunogen Information

Gene ID

18613

Swiss Prot

Q08481

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

Cd31; Pecam; PECAM-1

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification





Affinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

Validation Data



Flow cytometry: 1×10^6 BALB/c mouse splenocytes were surface-stained with Biotin Rabbit IgG isotype control (A25626, 5 µl/Test, left) or Biotin Rabbit anti-Mouse CD31 mAb (A27363, 0.25 µg, right), followed by APC Streptavidin staining.

Flow cytometry: 1×10^6 Rat splenocytes were surface-stained with Biotin Rabbit IgG isotype control (A25626, 5 µl/Test, left) or Biotin Rabbit anti-Mouse CD31 mAb (A27363, 0.5 µg, right), followed by APC Streptavidin staining.