PE Rabbit anti-Human CD2 mAb

Catalog No.: A27831



Basic Information

Observed MW

Calculated MW

39kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Human

CloneNo number

ARC59333

Conjugate

PE. Ex:565nm. Em:574nm.

Background

The protein encoded by this gene is a surface antigen found on all peripheral blood T-cells. The encoded protein interacts with LFA3 (CD58) on antigen presenting cells to optimize immune recognition. A locus control region (LCR) has been found in the 3' flanking sequence of this gene.

Recommended Dilutions

FC

5 μ l per 10^6 cells in 100 μ l volume

Immunogen Information

Gene ID 914 **Swiss Prot**

P06729

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

T11; SRBC; LFA-2

Contact

<u>a</u>		400-999-6126
\bowtie		cn.market@abclonal.com.cn
\overline{a}	ī	www.ahclonal.com.cn

Product Information


SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

Validation Data











Flow cytometry: 1X10^6 K-562 cells (negative control,left) and Jurkat cells (right) were surface-stained with PE Rabbit anti-Human CD2 mAb (A27831,5 μ I/Test,orange line) or PE Rabbit IgG isotype control (A24172,5 μ I/Test,blue line). Non-fluorescently stained cells were used as blank control (red line).



Flow cytometry: $1X10^6$ Jurkat cells were surface-stained with PE Rabbit IgG isotype control (A24172,5 μ I/Test,left) or PE Rabbit anti-Human CD2 mAb (A27831,5 μ I/Test,right).



Flow cytometry: 1X10^6 Human PBMC were surface-stained with PE Rabbit anti-Human CD2 mAb (A27831,5 µl/Test,orange line) or PE Rabbit IgG isotype control (A24172,5 µl/Test,blue line).Non-fluorescently stained cells were used as blank control (red line). Cells in the lymphocyte gate were used for analysis.

Flow cytometry: 1X10^6 Human PBMC were surface-stained with PE Rabbit IgG isotype control (A24172,5 μ I/Test,Ieft) or PE Rabbit anti-Human CD2 mAb (A27831,5 μ I/Test,right). Cells in the lymphocyte gate were used for analysis.