ABclonal www.abclonal.com

ABflo® 610 Rabbit anti-Human/Monkey CD64 mAb

Catalog No.: A27882

Basic Information

Observed MW

Calculated MW

43kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Human, Cynomolgus

CloneNo number

ARC53680

Conjugate

ABflo® 610. Ex:421nm. Em:612nm.

Background

This gene encodes a protein that plays an important role in the immune response. This protein is a high-affinity Fc-gamma receptor. The gene is one of three related gene family members located on chromosome 1.

Recommended Dilutions

FC

5 μ l per 10^6 cells in 100 μ l volume

Immunogen Information

Gene ID2209

Swiss Prot
P12314

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

CD64; FCG1; FCRI; CD64A; FCGR1; IGFR1; FcgammaRI

Contact

| 2 | | 400-999-6126 |
|-----------|---|---------------------------|
| \bowtie | | cn.market@abclonal.com.cn |
| • | T | www.abclonal.com.cn |

Product Information


SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.


Validation Data











Flow cytometry: 1X10^6 K-562 cells (negative control,left) and Human PBMC (right) were surface-stained with ABflo® 610 Rabbit anti-Human/Monkey CD64 mAb (A27882,5 µl/Test,orange line) or ABflo® 610 Rabbit IgG isotype control (A25826,5 µl/Test,blue line). Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry:1X10^6 Human PBMC were surface-stained with ABflo® 610 Rabbit IgG isotype control (A25826,5 μ I/Test,Ieft) or ABflo® 610 Rabbit anti-Human/Monkey CD64 mAb (A27882,5 μ I/Test,right). Cells in the monocyte gate were used for analysis.

Flow cytometry:1X10^6 Human PBMC were surface-stained with ABflo® 647 Rabbit anti-Human CD14 mAb (5 µl/Test) and ABflo® 610 Rabbit IgG isotype control (A25826,5 µl/Test,left) or ABflo® 610 Rabbit anti-Human/Monkey CD64 mAb (A27882,5 µl/Test,right). Cells in the lymphocyte and monocyte gates were used for analysis.