PTPN23 Rabbit pAb

Catalog No.: A4595



Basic Information

Observed MW

179kDa

Calculated MW

179kDa

Category

Primary antibody

Applications

WB,IHC-P,ELISA

Cross-Reactivity

Mouse, Rat

Background

This gene encodes a member of the non-receptor type protein-tyrosine phosphatase family. The encoded protein may be involved in the regulation of small nuclear ribonucleo protein assembly and pre-mRNA splicing by modifying the survival motor neuron (SMN) complex. The encoded protein additionally plays a role in ciliogenesis and is part of endosomal sorting complex required for transport (ESCRT) pathways. This gene may serve a tumor suppressor function.

Recommended Dilutions

WB 1:500 - 1:2000

IHC-P 1:50 - 1:200

Recommended starting concentration is 1 µg/mL.

Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene IDSwiss Prot
25930
Q9H3S7

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

HDPTP; HD-PTP; NEDBASS; PTP-TD14; PTPN23

Contact

6		400-999-6126
\bowtie		cn.market@abclonal.com.cn
<u> </u>	ī	www.ahclonal.com.cn


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.




Western blot analysis of various lysates using PTPN23 Rabbit pAb (A4595) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit lgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: $25\mu g$ per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 5s.



Immunohistochemistry analysis of paraffinembedded Rat brain using PTPN23 Rabbit pAb (A4595) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffinembedded Mouse brain using PTPN23 Rabbit pAb (A4595) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.