# TWIST2 Rabbit pAb

Catalog No.: A5599 1 Publications



# **Basic Information**

### **Observed MW**

18kDa

### **Calculated MW**

18kDa

### Category

Primary antibody

## **Applications**

WB,ELISA

#### **Cross-Reactivity**

Mouse, Rat

# **Background**

The protein encoded by this gene is a basic helix-loop-helix type transcription factor and shares similarity with Twist. This protein may inhibit osteoblast maturation and maintain cells in a preosteoblast phenotype during osteoblast development. This gene may be upregulated in certain cancers. Mutations in this gene cause focal facial dermal dysplasia 3, Setleis type. Two transcript variants encoding the same protein have been found.

# **Recommended Dilutions**

**WB** 1:500 - 1:1000

**ELISA** 

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

# **Immunogen Information**

 Gene ID
 Swiss Prot

 117581
 Q8WV|9

#### **Immunogen**

Synthetic peptide. This information is considered to be commercially sensitive.

## **Synonyms**

AMS; FFDD3; BBRSAY; DERMO1; SETLSS; bHLHa39; TWIST2

# **Contact**

<b>a</b>	400-999-6126
$\sim$	cn.market@abclonal.com.cn
<b>⊙</b>	www.abclonal.com.cn


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.



Western blot analysis of lysates from Rat uterus, using TWIST2 Rabbit pAb (A5599) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021).

Exposure time: 180s.