

TTC11/FIS1 Rabbit pAb

Catalog No.: A5821 **19 Publications**

Basic Information

Observed MW

17kDa

Calculated MW

17kDa

Category

Primary antibody

Applications

WB,IF/ICC,IP,ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

Enables identical protein binding activity. Involved in several processes, including calcium-mediated signaling using intracellular calcium source; cellular calcium ion homeostasis; and mitochondrion organization. Acts upstream of or within mitochondrion morphogenesis. Located in mitochondrion and peroxisome. Is integral component of mitochondrial outer membrane and integral component of peroxisomal membrane. Part of protein-containing complex. Biomarker of Alzheimer's disease.

Recommended Dilutions

WB 1:500 - 1:2000

IF/ICC 1:50 - 1:200

IP 0.5µg-4µg antibody for
200µg-400µg extracts of
whole cells

ELISA Recommended starting
concentration is 1 µg/mL.
Please optimize the
concentration based on
your specific assay
requirements.

Immunogen Information

Gene ID

51024

Swiss Prot

Q9Y3D6

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

TTC11; CGI-135; TTC11/FIS1

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.09% Sodium azide, 50% glycerol, pH7.3.


Validation Data



Immunoprecipitation analysis of 200 µg extracts of MCF7 cells using 3 µg TTC11/FIS1 antibody (A5821). Western blot was performed from the immunoprecipitate using TTC11/FIS1 antibody (A5821) at a dilution of 1:1000.



Western blot analysis of lysates from 293T cells, using TTC11/FIS1 Rabbit pAb (A5821) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 30s.



Confocal immunofluorescence analysis of HeLa cells using TTC11/FIS1 Rabbit pAb (A5821) at dilution of 1:50 (60x lens). Blue: DAPI for nuclear staining.