# **RICTOR Rabbit pAb**

Catalog No.: A6131 1 Publications



### **Basic Information**

#### **Observed MW**

200kDa

### **Calculated MW**

192kDa

#### Category

Primary antibody

### **Applications**

WB,ELISA

#### **Cross-Reactivity**

Rat

# **Background**

RICTOR and MTOR (FRAP1; MIM 601231) are components of a protein complex that integrates nutrient- and growth factor-derived signals to regulate cell growth (Sarbassov et al., 2004 [PubMed 15268862]).

## **Recommended Dilutions**

**WB** 1:500 - 1:1000

**ELISA** 

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

# **Immunogen Information**

**Gene ID**253260

Swiss Prot
Q6R327

#### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

### **Synonyms**

PIA; AVO3; hAVO3; RICTOR

### **Contact**

<b>a</b>	400-999-6126
×	cn.market@abclonal.com.cn
$\overline{a}$	www.ahclonal.com.cn

### **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

### **Storage**

Store at -20  $^{\circ}\text{C}.$  Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of lysates from PC-12 cells, using RICTOR Rabbit pAb (A6131) at 1:500 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 180s.