# Phospho-FLT3-Y589/591 Rabbit pAb

www.abclonal.com

**ABclonal** 

Catalog No.: AP0529 1 Publications

## **Basic Information**

#### **Observed MW**

160kDa

#### **Calculated MW**

113kDa

### Category

Primary antibody

## **Applications**

WB,ELISA

#### **Cross-Reactivity**

Human

# **Background**

This gene encodes a class III receptor tyrosine kinase that regulates hematopoiesis. This receptor is activated by binding of the fms-related tyrosine kinase 3 ligand to the extracellular domain, which induces homodimer formation in the plasma membrane leading to autophosphorylation of the receptor. The activated receptor kinase subsequently phosphorylates and activates multiple cytoplasmic effector molecules in pathways involved in apoptosis, proliferation, and differentiation of hematopoietic cells in bone marrow. Mutations that result in the constitutive activation of this receptor result in acute myeloid leukemia and acute lymphoblastic leukemia.

# **Recommended Dilutions**

**WB** 1:100 - 1:500

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

# **Immunogen Information**

**Gene ID Swiss Prot** 2322 P36888

#### **Immunogen**

Synthetic peptide. This information is considered to be commercially sensitive.

## **Synonyms**

FLK2; STK1; CD135; FLK-2; Phospho-FLT3-Y589/591

# **Contact**

6	400-999-6126
$\bowtie$	cn.market@abclonal.com.cr
$\odot$	www.abclonal.com.cr

## **Product Information**


Source Isotype **Purification** Rabbit IgG Affinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of lysates from A-431 cells, using Phospho-FLT3-Y589/591 Rabbit pAb (AP0529) at 1:500 dilution. A-431 cells were treated with pervanadate (1 mM) at 37°C for 30 minutes after serumstarvation overnight.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.