Leader in Biomolecular Solutions for Life Science

www.abclonal.com

Phospho-AKT1-T450 + AKT2-T451 + AKT3-T447 Rabbit mAb

Catalog No.: AP1430 Recombinant 1 Publications

Basic Information

Observed MW
60kDa

Calculated MW
48KDa/51KDa/55KDa

Category
Primary antibody

Applications
WB,IHC-P,ELISA

Cross-Reactivity
Human, Mouse, Rat

CloneNo number
ARC61098

Background

The serine-threonine protein kinase encoded by the AKT1 gene is catalytically inactive in serum-starved primary and immortalized fibroblasts. AKT1 and the related AKT2 are activated by platelet-derived growth factor. The activation is rapid and specific, and it is abrogated by mutations in the pleckstrin homology domain of AKT1. It was shown that the activation occurs through phosphatidylinositol 3-kinase. In the developing nervous system AKT is a critical mediator of growth factor-induced neuronal survival. Survival factors can suppress apoptosis in a transcription-independent manner by activating the serine/threonine kinase AKT1, which then phosphorylates and inactivates components of the apoptotic machinery. Mutations in this gene have been associated with the Proteus syndrome. Multiple alternatively spliced transcript variants have been found for this gene.

Recommended Dilutions

WB	1:2000 - 1:10000
IHC-P	1:100 - 1:500
ELISA	Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID
207/208/10000

Swiss Prot
P31749/P31751/Q9Y243

Immunogen
Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

Contact

400-999-6126

cn.market@abclonal.com.cn

www.abclonal.com.cn

Product Information

Source
Rabbit

Isotype
IgG

Purification
Affinity purification

Storage
Store at -20℃. Avoid freeze / thaw cycles.
Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Western blot analysis of lysates from RD cells, using Phospho-AKT1-T450 + AKT2-T451 + AKT3-T447 Rabbit mAb (AP1430) at 1:10000 dilution. RD cells were treated by λ-PP mixed solution (1ul) at 30℃ for 30 minutes.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 30s.



Western blot analysis of lysates from C2C12 cells, using Phospho-AKT1-T450 + AKT2-T451 + AKT3-T447 Rabbit mAb (AP1430) at 1:10000 dilution. C2C12 cells were treated by λ-PP mixed solution (1ul) at 30℃ for 30 minutes.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 30s.



Immunohistochemistry analysis of paraffin-embedded Human breast tissue using Phospho-AKT1-T450 + AKT2-T451 + AKT3-T447 Rabbit mAb (AP1430) at a dilution of 1:300 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human tonsil tissue using Phospho-AKT1-T450 + AKT2-T451 + AKT3-T447 Rabbit mAb (AP1430) at a dilution of 1:300 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse intestin tissue using Phospho-AKT1-T450 + AKT2-T451 + AKT3-T447 Rabbit mAb (AP1430) at a dilution of 1:300 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.

Validation Data



Immunohistochemistry analysis of paraffin-embedded Mouse kidney tissue using Phospho-AKT1-T450 + AKT2-T451 + AKT3-T447 Rabbit mAb (AP1430) at a dilution of 1:300 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat intestine tissue using Phospho-AKT1-T450 + AKT2-T451 + AKT3-T447 Rabbit mAb (AP1430) at a dilution of 1:300 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat kidney tissue using Phospho-AKT1-T450 + AKT2-T451 + AKT3-T447 Rabbit mAb (AP1430) at a dilution of 1:300 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.

For research use only. Not for therapeutic or diagnostic purposes.
Please visit http://abclonal.com for a complete listing of recommended products.