

# Phospho-Akt-S473 Rabbit mAb

Catalog No.: AP1453   **Recombinant**   **6 Publications**

## Basic Information

### Observed MW

60kDa

### Calculated MW

48kDa/55kDa/51kDa/54kDa

### Category

Primary antibody

### Applications

WB,IHC-P,ELISA

### Cross-Reactivity

Human, Mouse, Rat

### Clone/No. number

ARC5023-02

## Background

The serine-threonine protein kinase encoded by the AKT1 gene is catalytically inactive in serum-starved primary and immortalized fibroblasts. AKT1 and the related AKT2 are activated by platelet-derived growth factor. The activation is rapid and specific, and it is abrogated by mutations in the pleckstrin homology domain of AKT1. It was shown that the activation occurs through phosphatidylinositol 3-kinase. In the developing nervous system AKT is a critical mediator of growth factor-induced neuronal survival. Survival factors can suppress apoptosis in a transcription-independent manner by activating the serine/threonine kinase AKT1, which then phosphorylates and inactivates components of the apoptotic machinery. Mutations in this gene have been associated with the Proteus syndrome. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jul 2011]

## Recommended Dilutions

**WB**                    1:1000 - 1:3000

**IHC-P**                1:50 - 1:200

**ELISA**                Recommended starting concentration is 1  $\mu$ g/mL. Please optimize the concentration based on your specific assay requirements. For high-ratio antibody dilutions ( $\geq 1:10000$ ) a sequential dilution method is strongly recommended to ensure measurement accuracy.

## Immunogen Information

### Gene ID

207/208/10000

### Swiss Prot

P31749/P31751/Q9Y243

### Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

### Synonyms

AKT1/AKT2/AKT3; Phospho-Akt-S473

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification

Affinity purification

### Storage



Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


## Contact

	400-999-6126
	<a href="mailto:cn.market@abclonal.com.cn">cn.market@abclonal.com.cn</a>
	<a href="http://www.abclonal.com.cn">www.abclonal.com.cn</a>


## Validation Data




Immunohistochemistry analysis of paraffin-embedded Mouse brain tissue using Phospho-Akt-S473 Rabbit mAb (AP1453) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse colon tissue using Phospho-Akt-S473 Rabbit mAb (AP1453) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse testis tissue using Phospho-Akt-S473 Rabbit mAb (AP1453) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat kidney tissue using Phospho-Akt-S473 Rabbit mAb (AP1453) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat intestine tissue, Untreated (left) and lambda phosphatase-treated (right), using Phospho-Akt-S473 Rabbit mAb (AP1453) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse spleen tissue, Untreated (left) and lambda phosphatase-treated (right), using Phospho-Akt-S473 Rabbit mAb (AP1453) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.