

# ENO1 Knockdown HeLa Cell Line, Heterozygous

**Catalog No.:** RM01838

## Basic Information

**Catalog No.**

RM01838

**Category**

Cell Line

**Parental Cell line**

HeLa

**Genotype**

Knockdown

## Gene Information

**Gene Symbol**

ENO1

**Species**

Human

**Gene ID**

2023

**Swiss Prot**

P06733

**Synonyms**

ENO1L1; HEL-S-17; MPB1; NNE; PPH

## Contact

 | 400-999-6126 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn) | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Background

This gene encodes alpha-enolase, one of three enolase isoenzymes found in mammals. Each isoenzyme is a homodimer composed of 2 alpha, 2 gamma, or 2 beta subunits, and functions as a glycolytic enzyme. Alpha-enolase in addition, functions as a structural lens protein (tau-crystallin) in the monomeric form. Alternative splicing of this gene results in a shorter isoform that has been shown to bind to the c-myc promoter and function as a tumor suppressor. Several pseudogenes have been identified, including one on the long arm of chromosome 1. Alpha-enolase has also been identified as an autoantigen in Hashimoto encephalopathy.

## Product Information

**Description**

ENO1 Knockdown HeLa cell line is engineered from HeLa cell line with Gene-Editing Technology.

Allele-1:WT

Allele-2:exon3 was deleted

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

**Packaging**

1 vial parental cell line and 1 vial knockout cell line

**Shipping Conditions**

Dry ice

**Amount**

1~5x10<sup>6</sup> cells/vial.

**Storage**

Stored in liquid nitrogen for a long time less than -130°C. Minimizing freeze-thaw cycles.

**Protocol**

Upon arrival, it should be maintained in DMEM medium with 10%(v/v) fetal bovine serum and 100U penicillin-streptomycin, at 37°C with 5% CO<sub>2</sub> condition.


1. Thaw the vial in 37°C water bath, and shake it to melt as soon as possible.
2. Transfer the cell suspension to a 15mL conical tube with pre-warmed 5mL complete medium and centrifuge 1000rpm for approximately 5 minutes at room temperature.
3. Remove and discard the supernatant.
4. Resuspend the cell pellet with 1mL pre-warmed complete medium and seed in 10cm dish.
5. Add 8-10mL of complete medium.
6. Incubate the culture at 37°C incubator with 5% CO<sub>2</sub>.
7. A subcultivation ratio of 1:2-1:4 is recommended.

## Sequencing data

WT CAGTGGTTCTCTCT\*\*\*\*\*TAATGCCACCAGAG  
Mut CAGTGGTTCTCTCT\*\*\*\*\*TAATGCCACCAGAG  
Allele-1: WT  
WT CGCGTCGGCCTCAA\*\*\*\*\*TCCCAGGCCAGGG  
Mut CGCGTCGGCCTCAA\*\*\*Deletion\*\*\*TCCCAGGCCAGGG  
Allele-2: exon3 was deleted

Genome sequence analysis of PCR products from parental (WT) and ENO1 Knockdown (KD) HeLa cells, using sanger sequencing.

## WB data



Western blot analysis of extracts from parental (Control) and ENO1 knockdown (KD) HeLa cells, using ENO1 antibody at 1:3000 dilution.