

KDR Knockout HeLa Cell Lysate, Homozygous

Catalog No.: RM02021

Basic Information

Catalog No.

RM02021

Category

Cell Lysate

Parental Cell line

HeLa

Genotype

Knockout

Gene Information

Gene Symbol

KDR

Species

Human

Gene ID

3791

Swiss Prot

P35968

Synonyms

CD309; FLK1; VEGFR; VEGFR2

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Background

Vascular endothelial growth factor (VEGF) is a major growth factor for endothelial cells. This gene encodes one of the two receptors of the VEGF. This receptor, known as kinase insert domain receptor, is a type III receptor tyrosine kinase. It functions as the main mediator of VEGF-induced endothelial proliferation, survival, migration, tubular morphogenesis and sprouting. The signalling and trafficking of this receptor are regulated by multiple factors, including Rab GTPase, P2Y purine nucleotide receptor, integrin alphaVbeta3, T-cell protein tyrosine phosphatase, etc.. Mutations of this gene are implicated in infantile capillary hemangiomas. [provided by RefSeq, May 2009]

Product Information

Description

KDR Knockout HeLa Cell Line is engineered from HeLa cell line with Gene-Editing technology.

Allele-1:1bp deletion in exon3

Allele-2:1bp deletion in exon3

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles.

Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins.

Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell Lysate and 1 vial knockout cell Lysate

Shipping Conditions

4°C

Amount

50μL, 2μg/μL.

Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

Protocol

To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

Sequencing data

WT GTGGAGGTGACTGAGTGCAGCGATGGCCTCTTCTGTAA
Mut GTGGAGGTGACTGAGTGCA -CGATGGCCTCTTCTGTAA
Allele-1: 1bp deletion in exon3

WT GTGGAGGTGACTGAGTGCAGCGATGGCCTCTTCTGTAA
Mut GTGGAGGTGACTGAGTGCA -CGATGGCCTCTTCTGTAA
Allele-2: 1bp deletion in exon3

Genome sequence analysis of PCR products from parental (WT) and KDR knockout (KO) HeLa cells, using sanger sequencing.