

# MAP3K8 Knockout 293T Cell Line, Homozygous

**Catalog No.:** RM02187

## Basic Information

**Catalog No.**

RM02187

**Category**

Cell Line

**Parental Cell line**

293T

**Genotype**

Knockout

## Gene Information

**Gene Symbol**

MAP3K8

**Species**

Human

**Gene ID**

1326

**Swiss Prot**

P41279

**Synonyms**AURA2; COT; EST; ESTF; MEKK8; TPL2;  
Tpl-2; c-COT

## Contact

 | 400-999-6126 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn) | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Background

This gene is an oncogene that encodes a member of the serine/threonine protein kinase family. The encoded protein localizes to the cytoplasm and can activate both the MAP kinase and JNK kinase pathways. This protein was shown to activate I $\kappa$ B kinases, and thus induce the nuclear production of NF- $\kappa$ B. This protein was also found to promote the production of TNF- $\alpha$  and IL-2 during T lymphocyte activation. This gene may also utilize a downstream in-frame translation start codon, and thus produce an isoform containing a shorter N-terminus. The shorter isoform has been shown to display weaker transforming activity. Alternate splicing results in multiple transcript variants that encode the same protein. [provided by RefSeq, Sep 2011]

## Product Information

**Description**

MAP3K8 Knockout 293T Cell Line is engineered from 293T cell line with Gene-Editing Technology.

Allele-1:54bp insertion and 1bp deletion in exon2

Allele-2:58bp deletion in exon2

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

**Packaging**

1 vial parental cell line and 1 vial knockout cell line

**Shipping Conditions**

Dry ice

**Amount**

1~5x10<sup>6</sup> cells/vial

**Storage**

Stored in liquid nitrogen for a long time less than -130°C. Minimizing freeze-thaw cycles.

**Protocol**

Upon arrival, it should be maintained in DMEM medium with 10%(v/v) fetal bovine serum and 100U penicillin-streptomycin, at 37°C with 5% CO<sub>2</sub> condition.

1. Thaw the vial in 37°C water bath, and shake it to melt as soon as possible.
2. Transfer the cell suspension to a 15mL conical tube with pre-warmed 5mL complete medium and centrifuge 1000rpm for approximately 5 minutes at room temperature.
3. Remove and discard the supernatant.
4. Resuspend the cell pellet with 1mL pre-warmed complete medium and seed in 10cm dish.
5. Add 8-10mL of complete medium.
6. Incubate the culture at 37°C incubator with 5% CO<sub>2</sub>.
7. A subcultivation ratio of 1:2-1:4 is recommended.

# Sequencing data

WT ATGTTCTCCTGATC\*\*\*\*\*CCCTGGAAGCTGAC\*\*TCGGGGCGCCTTTGGAAAGGTATACTTGG  
Mut ATGTTCTCCTGATC\*\*Insertion\*\*CCCTGGAAGCTGAC\*\*TCGGGGCGCCTTTG AAAGGTATACTTGG  
Allele-1: 54bp insertion and 1bp deletion in exon2  
WT ATGTTCTCCTGATC\*\*\*\*\*GAAAGGTATACTTG  
Mut ATGTTCTCCTGATC\*\*Deletion\*\*GAAAGGTATACTTG  
Allele-2: 58bp deletion in exon2

Genome sequence analysis of PCR products from parental (WT) and MAP3K8 knockout (KO) 293T cells, using sanger sequencing.