

# SLC25A4 Knockout 293T Cell Lysate, Homozygous

Catalog No.: RM02293

#### **Basic Information**

#### Catalog No.

RM02293

#### Category

Cell Lysate

#### **Parental Cell line**

293T

#### Genotype

Knockout

## **Background**

This gene is a member of the mitochondrial carrier subfamily of solute carrier protein genes. The product of this gene functions as a gated pore that translocates ADP from the cytoplasm into the mitochondrial matrix and ATP from the mitochondrial matrix into the cytoplasm. The protein forms a homodimer embedded in the inner mitochondria membrane. Mutations in this gene have been shown to result in autosomal dominant progressive external opthalmoplegia and familial hypertrophic cardiomyopathy. [provided by RefSeq, Jun 2013]

#### **Gene Information**

#### **Gene Symbol**

SLC25A4

#### **Species**

Human

#### **Gene ID**

291

#### **Swiss Prot**

P12235

#### **Synonyms**

AAC1; ANT; ANT 1; ANT1; MTDPS12; MTDPS12A; PEO2; PEO3; PEOA2; T1

### **Contact**

<b>a</b>	400-999-6126
$\bowtie$	cn.market@abclonal.com.cn
•	www.abclonal.com.cn

#### **Product Information**

#### Description

SLC25A4 Knockout 293T Cell Line is engineered from 293T cell line with Gene-Editing technology.

Allele-1:124bp deletion in exon2

Allele-2:124bp deletion in exon2

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

#### **Packaging**

 ${\bf 1}$  vial parental cell Lysate and  ${\bf 1}$  vial knockout cell Lysate

# **Shipping Conditions**

Amount

4°C

50μL, 2μg/μL.

### Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

#### Protoco

To be used as WB control. Lysate is supplied in  $1\times$  SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

# Sequencing data

WT TAACCTGGCCAACG\*CGGTGGGGCCGCTG
Mut TAACCTGGCCAACG\*\*\*Deletion\*\*\*CGGTGGGGCCGCTG
Allele-1: 124bp deletion in exon2

WT TAACCTGGCCAACG\*CGGTGGGGCCGCTG
Mut TAACCTGGCCAACG\*\*\*\*Deletion\*\*\*\*CGGTGGGGCCGCTG

Allele-2: 124bp deletion in exon2

Genome sequence analysis of PCR products from parental (WT) and SLC25A4 knockout (KO) 293T cells, using sanger sequencing.