

# Recombinant Human IL-11 Protein

Catalog No.: RP00050    Recombinant    1 Publications

## Sequence Information

**Species**    **Gene ID**    **Swiss Prot**  
 Human        3589        P20809

### Tags

No tag

### Synonyms

IL11;AGIF;IL-11

## Background

The protein is a member of the gp130 family of cytokines. These cytokines drive the assembly of multisubunit receptor complexes, all of which contain at least one molecule of the transmembrane signaling receptor IL6ST (gp130). This cytokine is shown to stimulate the T-cell-dependent development of immunoglobulin-producing B cells. It is also found to support the proliferation of hematopoietic stem cells and megakaryocyte progenitor cells.

## Basic Information

### Description

Recombinant Human IL-11 Protein is produced by *E. coli* expression system. The target protein is expressed with sequence (Pro22-Leu199) of human IL-11 (Accession #NP\_000632.1) fused with an initial Met at the N-terminus.

### Bio-Activity

Measured in a cell proliferation assay using TF-1 human erythroleukemic cells. The ED<sub>50</sub> for this effect is 1.3-5 ng/mL, corresponding to a specific activity of 2×10<sup>5</sup>-7.69×10<sup>5</sup>units/mg.

### Shipping

The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

### Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

### Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for handling.

## Product Information

**Source**    **Purification**  
*E. coli*        ≥ 95 % as  
 determined by SDS-PAGE.

**Calculated MW**    **Observed MW**  
 19.14 kDa        21 kDa

**Endotoxin**  
 < 1 EU/μg of the protein by LAL method.

**Formulation**  
 Lyophilized from a 0.22 μm filtered solution of 20mM Tris, 150mM NaCl, pH 8.0. Contact us for customized product form or formulation.


**Reconstitution**  
 Centrifuge the vial before opening.

Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize freeze-thaw cycles.


## Contact

	400-999-6126
	<a href="mailto:cn.market@abclonal.com.cn">cn.market@abclonal.com.cn</a>
	<a href="http://www.abclonal.com.cn">www.abclonal.com.cn</a>

## Validation Data



Recombinant Human IL-11 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



Recombinant human IL11 promotes the proliferation of TF-1 human erythroleukemic cells. The ED<sub>50</sub> for this effect is 1.3-5ng/mL, corresponding to a specific activity of  $2 \times 10^5$ - $7.69 \times 10^5$ units/mg.