

Catalog No.: RP01771 **Recombinant**

Species	Gene ID	Swiss Prot
Mouse	50706	O62009-2

C-His

PN; PLF; Osf2; OSF-2;
A630052E07Bik□POSTN

Source	Purification
HEK293 cells	≥ 95% as determined by SDS-PAGE.

70.64 kDa 70-75 kDa

< 0.1 EU/μg of the protein by LAL method.

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Periostin (POSTN), also known as OSF2 (osteoblast specific factor 2), is a heterofunctional secreted extracellular matrix (ECM) protein comprised of four fasciclin domains that promotes cellular adhesion and movement, as well as collagen fibrillogenesis. Postn is frequently overexpressed in various types of human cancers, stimulating metastatic growth by promoting cancer cell survival, invasion and angiogenesis, and can be a useful marker to predict the behavior of cancer.

Recombinant Mouse Periostin/PN/POSTN Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Asn24-Ile653) of mouse Periostin/PN/POSTN (Accession #NP_056599.1) fused with a 6×His tag at the C-terminus.

Measured by its ability to induce adhesion of ATDC5 mouse chondrogenic cells. Immobilized Recombinant Human Periostin/OSF-2 at 10 µg/mL (100 µL/well) induces >13% cell adhesion. Measured by its ability to induce adhesion of ATDC5 mouse chondrogenic cells.


Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.
After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.
Avoid repeated freeze/thaw cycles.

 | 400-999-6126

✉ | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Validation Data



Recombinant Mouse Periostin/PN/POSTN Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.