

Recombinant Rat Hepatocyte growth factor/HGF Protein

Catalog No.: RP01776 **Recombinant**

Sequence Information

Species **Gene ID** **Swiss Prot**
 Rat 24446 P17945

Tags

C-His

Synonyms

HPTA|Hepatocyte growth factor|HGF

Background

Hepatocyte growth factor/scatter factor (HGF/SF) is a paracrine cellular growth, motility and morphogenic factor. It belongs to the peptidase S1 family and Plasminogen subfamily, contains 4 kringle domains, 1 PAN domain and 1 peptidase S1 domain. HGF regulates cell growth, cell motility, and morphogenesis by activating tyrosine kinase signaling cascade after binding to the proto-oncogenic c-Met receptor. HGF is secreted by mesenchymal cells and acts as a multi-functional cytokine on cells of mainly epithelial origin. Its ability to stimulate mitogenesis, cell motility, and matrix invasion gives it a central role in angiogenesis, tumorogenesis, and tissue regeneration.

Basic Information

Description

Recombinant Rat Hepatocyte growth factor/HGF Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Gln 33-Leu728) of rat HGF (Accession #NP_058713.1) fused with and a 6xHis tag at the C-terminus.

Bio-Activity

Measured in a cell proliferation assay using Mv.1.lu cells. The ED₅₀ for this effect is 1.6-6.2 ng/mL, corresponding to a specific activity of 1.6×10⁵~6.3×10⁵ units/mg.

Shipping

The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for handling.

Product Information

Source **Purification**
 HEK293 cells ≥ 90 % as
 determined by SDS-PAGE.

Calculated MW **Observed MW**
 80.14 kDa 80-100 kDa

Endotoxin

< 0.1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.


Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize freeze-thaw cycles.

Contact

	400-999-6126
	cn.market@abclonal.com.cn
	www.abclonal.com.cn

Validation Data



Recombinant Rat Hepatocyte growth factor/HGF Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.

Recombinant Rat HGF stimulates cell proliferation of the Mv.1.lu cells. The ED₅₀ for this effect is 1.6-6.2 ng/mL, corresponding to a specific activity of 1.6×10⁵~6.3×10⁵ units/mg.