

Catalog No.: RP01797 **Recombinant**

Species	Gene ID	Swiss Prot
Human	2324	P35916

C-His

PCL; CHTD7; FLT-4; FLT41; LMPH1A;
LMPHM1; VEGFR3; VEGFR-3

Source	Purification
HEK293 cells	≥ 90 % as determined by SDS-PAGE.

Calculated MW	Observed MW
85.37 kDa	60, 70, 140 kDa

< 0.01 EU/μg of the protein by LAL method

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

 | 400-999-6126

✉ | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Vascular endothelial growth factor receptor 3 (VEGFR3), also known as FLT-4, together with the other two members VEGFR1 (FLT-1) and VEGFR2 (KDR/Flk-1) are receptors for vascular endothelial growth factors (VEGF) and belong to the class III subfamily of receptor tyrosine kinases (RTKs). The VEGFR3 protein is expressed mainly on lymphatic vessels but it is also up-regulated in tumor angiogenesis. Mutations in VEGFR3 have been identified in patients with primary lymphoedema. The VEGF-C/VEGF-D/VEGFR3 signaling pathway may provide a target for antilymphangiogenic therapy in prostate cancer, breast cancer, gastric cancer, lung cancer, non-small cell lung cancer (NSCLC), and so on.


Recombinant Human VEGFR-3/FLT-4 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Tyr25-Ile776) of human VEGFR-3/FLT-4 (Accession #NP_002011.2) fused with 6xHis tag at the C-terminus.

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human VEGFR-3/FLT-4 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.