

Catalog No.: RP01813 **Recombinant**

Species	Gene ID	Swiss Prot
Human	4681	P41271-1

C-hFc

NB: DAN; NO3; DAND1; D1S1733E

Source	Purification
HEK293 cells	≥ 95 % as determined by SDS-PAGE

Calculated MW	Observed MW
43.66 kDa	60 kDa

< 0.01 EU/μg of the protein by LAL method

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

 | 400-999-6126

 | cn.market@abclonal.com.cn


 | www.abclonal.com.cn

The Dan (Differential screening-selected gene aberrative in neuroblastoma, also known as N03) gene was first identified as the putative rat tumor suppressor gene and encodes a protein structurally related to Cerberus and Gremlin in the vertebrates. It is a founding member of the DAN family of secreted proteins, acts as an inhibitor of cell cycle progression, and is closely involved in retinoic acid-induced neuroblastoma differentiation. There are at least five mammalian protein members in the evolutionarily conserved Dan family including DAN, Gremlin/DRM, Cer1 (Cerberus-related), Dante, and PRDC (protein related to DAN and Cerberus), and share the C-terminal cystine-knot motif. As a secreted glycoprotein, DAN is a member of a class of glycoproteins shown to be secreted inhibitors of the transforming growth factor-beta (TGF-beta) and bone morphogenic protein pathways. It binds to BMPs and preventing their interactions with signaling receptor complexes, and accordingly regulates the processes of embryonic development and tissue differentiation. DAN gene product may have an important role in the regulation of the entry of cells into the S phase. Besides, the DAN gene product possesses an ability to revert phenotypes of transformed rat fibroblasts and represents a candidate tumor suppressor gene for neuroblastoma.

Recombinant Human NBL1 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Ala17-Asp181) of human NBL1 (Accession #NP_002183.1) fused with a hFc tag at the C-terminus.

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.
After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.
Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human NBL1 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.