

Recombinant Human CCL14/HCC-1/HCC-3(28-93) Protein

Catalog No.: RP01985 Recombinant

Sequence Information

Species Gene ID Swiss Prot Human 6358(9-74) Q16627

Tags C-6His

Synonyms

CCL14; NCC2; SCYA14;C-C motif chemokine 14; Chemokine CC-1/CC-3; HCC-1/HCC-3; HCC-1(1-74); NCC-2; Small-inducible cytokine A14; Cleaved into: HCC-1(3-74); HCC-1(4-74); HCC-1(9-74)

Product Information

Source Purification HEK293 cells ≥ 95 % as

determined by SDS-

PAGE.

Calculated MW Observed MW

8.623 KDa 10-15 kDa

Endotoxin

< 0.1 EU/µg of the protein by LAL method.

Formulation

Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

Background

Chemokine (C-C motif) Ligand 14 (CCL14) is a small cytokine belonging to the CC chemokine family. It isproduced as a protein precursor that is processed to generate a mature active protein containing 74 aminoacids that and is 46% identical in amino acid composition to CCL3 and CCL4. This chemokine is expressed invarious tissues including spleen, bone marrow, liver, muscle, and gut. CCL14 activates monocytes, but does notinduce their chemotaxis. Human CCL14 is located on chromosome 17 within a cluster of other chemokinesbelonging to the CC family.

Basic Information

Description

Recombinant Human CCL14/HCC-1/HCC-3(28-93) Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Gly28-Asn93) of Human CCL14/HCC-1/HCC-3(28-93) (Accession #NP_116739.1) fused with His at the Cterminus.

Bio-Activity


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20 $^{\circ}$ C for 3 months, at 2-8 $^{\circ}$ C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human CCL14/HCC-1/HCC-3(28-93)Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.