# Recombinant Human SR-B3/CD36 Protein

ABclonal

www.abclonal.com

Catalog No.: RP02303 Recombinant

# **Sequence Information**

**Species** Gene ID **Swiss Prot** Human 948 P16671

**Tags** 

C-His

**Synonyms** 

BDPLT10;CHDS7;FAT;GP3B;GP4;GPIV;PA SIV;SCARB3;CD36; BDPLT10; CHDS7; FAT; GP3B; GP4; GPIV; PASIV; SCARB3; CD36 molecule

# **Product Information**

**Purification** HEK293 cells

≥ 95 % as determined by SDS-

PAGE.

Calculated MW Observed MW

47.46 kDa 75-100 kDa

#### Endotoxin

< 0.1 EU/µg of the protein by LAL method.

#### **Formulation**

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.

# Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

### Contact

<u>a</u>	400-999-6126
$\bowtie$	cn.market@abclonal.com.cn
•	www.abclonal.com.cn

# **Background**

Platelet Glycoprotein 4 (CD36) is an integral membrane glycoprotein that has multiple physiological functions. It is broadly expressed on a variety of cell types including microvascular endothelium, adipocytes, skeletal muscle, epithelial cells of the retina, breast, and intestine, smooth muscle cells, erythroid precursors, platelets, megakaryocytes, dendritic cells, monocytes/macrophages, and microglia. As a member of the scavenger receptor family, CD36 is a multiligand pattern recognition receptor that interacts with a large number of structurally dissimilar ligands, including long chain fatty acid (LCFA), advanced glycation end products (AGE), thrombospondin-1,oxidized lowdensity lipoproteins (oxLDLs), high density lipoprotein (HDL), phosphatidylserine, apoptotic cells,  $\beta$  amyloid fibrils (fA $\beta$ ), collagens I and IV, and Plasmodium falciparuminfected erythrocytes. CD36 is required for the antiangiogenic effects of thrombospondin-1 in the corneal neovascularization assay. It plays a role in lipid metabolism and has been identified as a fatty acid translocase necessary for the binding and transport of LCFA in cells and tissues.

## **Basic Information**

#### Description

Recombinant Human CD36 Protein is produced by Expi293 expression system. The target protein is expressed with sequence (Gly30-Asn439) of Human CD36 fused with a His tag at the C-terminal.

#### **Bio-Activity**


### Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

# **Validation Data**



Recombinant Human SR-B3/CD36 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.