

Biotinylated Recombinant Human Low-density lipoprotein receptor/LDLR Protein

Catalog No.: RP02430B Recombinant

Sequence Information

Species Gene ID Swiss Prot Human 3949 P01130

Tags C-His&Avi

Synonyms

LDLR; FH; FHC; LDLCQ2; low-density lipoprotein receptor; FH; FHC; LDLCQ2

Product Information

Source Purification

HEK293 cells ≥ 95 % as determined by Tris-Bis PAGE;≥ 95 % as determined by

HPLC.

Calculated MW Observed MW

87.6 kDa 120-140 kDa

Endotoxin

< 1 EU/ μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μ m filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

6	400-999-6126
\bowtie	cn.market@abclonal.com.cn
•	www.abclonal.com.cn

Background

The low density lipoprotein receptor (LDLR) is the founding member of the LDL R family of widely expressed cell surface scavenger receptors. It is a cell-surface receptor that recognizes the apoprotein B100 which is embedded in the phospholipid outer layer of LDL particles.

Basic Information

Description

Biotinylated Recombinant Human Low-density lipoprotein receptor/LDLR Protein is produced by Expi293 expression system. The target protein is expressed with sequence (Ala22-Arg788) of Human LDLR fused with a His tag and Avi tag at the Cterminal.

Bio-Activity


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80°C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Biotinylated Recombinant Human Lowdensity lipoprotein receptor/LDLR Protein was determined by Tris-Bis PAGE under reducing conditions with Coomassie Blue.



The purity of Biotinylated Human LDLR is greater than 95% as determined by SEC-HPLC.