

Recombinant Human CDK17&Cyclin Y Kinase

Catalog No.: RP03307LQ Recombinant

Sequence Information

 Species
 Gene ID
 Swiss Prot

 Human
 5128&21977
 Q00537&Q8

 1
 ND76

Tags

N-GST (CDK17) & N-Flag (Cyclin Y)

Synonyms

CDK17; PCTAIRE2; PCTK2; Cyclindependent kinase 17; Cyclin Y; CCNY; C10orf9; CBCP1; CFP1; Cyclin-Y

Product Information

Source	Purification
Baculovirus-Insect	≥ 90 % as
Cells	determined by SDS-
	PAGE;≥ 90 % as
	determined by
	HPLC.

Calculated MW Observed MW

86.1 kDa/41.3 kDa 75-85 kDa/25-40

kDa

Endotoxin

< 1 EU/µg of the protein by LAL method.

Formulation

Supplied as a 0.22 μ m filtered solution in 20 mM HEPES, 200 mM NaCl, 5% glycerol, 1 mM DTT. (pH 7.5). Contact us for customized product form or formulation.

Reconstitution

Please use running water to thaw it quickly.

Contact

<u>a</u>		400-999-6126
\bowtie		cn.market@abclonal.com.cn
•	T	www.abclonal.com.cn

Background

CDK17 also known as Serine/threonine-protein kinase PCTAIRE-2 belongs to the cdc2/cdkx subfamily of the ser/thr family of protein kinases. It is highly expressed in terminally differentiated neurons, particularly those found in hippocampal regions and olfactory bulbs.

Basic Information

Description

Recombinant Human CDK17&Cyclin Y Protein is produced by Baculovirus-Insect Cells expression system. The target protein is expressed with sequence (Met1-Phe523 (CDK17) & Met1-Ser341 (Cyclin Y)) of Human CDK17&CCNY (Accession #Q00537&Q8ND76) fused with a N-GST (CDK17) & N-Flag (Cyclin Y) tag.

Bio-Activity

The activity of CDK17/Cyclin Y is based on the MSA technology, and the content and ratio of the substrate and the product are directly separated and detected in real time and dynamically by the different migration rates of the substrate and the product after the enzymatic reaction.

Storage


Store at -70°C. This product is stable at \leq -70°C for up to 1 year from the date of receipt. For optimal storage, aliquot into smaller quantities after centrifugation and store at recommended temperature.

Aliquots below 10 μ L are not advisable. Product must not be stored in diluted solutions. Avoid repeated freeze-thaw cycles.

Avoid repeated freeze/thaw cycles.



Recombinant Human CDK17&Cyclin Y Kinase was determined by SDS-PAGE under reducing (R) and non-reducing (NR) conditions.



The activity of CDK17/Cyclin Y is based on the MSA technology, and the content and ratio of the substrate and the product are directly separated and detected in real time and dynamically by the different migration rates of the substrate and the product after the enzymatic reaction.